

PROLAPSE QUESTIONNAIRE

NAME: _____

PFDI- 20 Instructions: Please answer all of the questions in the following survey. These questions will ask you if you have certain bowel, bladder, or pelvic symptoms and, if you do, how much they bother you. Answer these by circling the appropriate number. While answering these questions, please consider your symptoms over the last 3 months.

The PFDI-20 has 20 items and 3 scales of your symptoms.

All items use the following format with a response scale from 0 to 4.

Symptoms Not Present = NO

0 = Not present

Symptoms Present = YES, scale of bother:

1 = not at all

2 = somewhat

3 = moderately

4 = quite a bit

Pelvic Organ Prolapse Distress Inventory 6 (POPDI-6)

Do you ...	No	Yes			
1. Usually experience pressure in the lower abdomen?	0	1	2	3	4
2. Usually experience heaviness or dullness in the pelvic area?	0	1	2	3	4
3. Usually have a bulge or something falling out that you can see or feel in your vaginal area?	0	1	2	3	4
4. Ever have to push on the vagina or around the rectum to have or complete a bowel movement?	0	1	2	3	4
5. Usually experience a feeling of incomplete bladder emptying?	0	1	2	3	4
6. Ever have to push up on a bulge in the vaginal area with your fingers to start or complete urination?	0	1	2	3	4

Colorectal-Anal Distress Inventory 8 (CRAD-8):

Do you ...	No	Yes			
7. Feel you need to strain too hard to have a bowel movement?	0	1	2	3	4
8. Feel you have not completely emptied your bowels at the end of a bowel movement?	0	1	2	3	4
9. Usually lose stool beyond your control if your stool is well formed?	0	1	2	3	4
10. Usually lose stool beyond your control if your stool is loose?	0	1	2	3	4
11. Usually lose gas from the rectum beyond your control?	0	1	2	3	4
12. Usually have pain when you pass your stool?	0	1	2	3	4
13. Experience a strong sense of urgency and have to rush to the bathroom to have a bowel movement?	0	1	2	3	4
14. Does part of your bowel ever pass through the rectum and bulge outside during or after a bowel movement?	0	1	2	3	4

Urinary Distress Inventory 6 (UDI-6):

Do you ...	No	Yes			
15. Usually experience frequent urination?	0	1	2	3	4
16. Usually experience urine leakage associated with a feeling of urgency, that is, a strong sensation of needing to go to the bathroom?	0	1	2	3	4
17. Usually experience urine leakage related to coughing, sneezing, or laughing?	0	1	2	3	4
18. Usually experience small amounts of urine leakage (that is, drops)?	0	1	2	3	4
19. Usually experience difficulty emptying your bladder?	0	1	2	3	4
20. Usually experience <i>pain</i> or <i>discomfort</i> in the lower abdomen or genital region?	0	1	2	3	4

Scoring the PFDI-20:

Scale Scores: Obtain the mean value of all of the answered items within the corresponding scale (possible value 0 to 4) and then multiply by 25 to obtain the scale score (range 0 to 100). Missing items are dealt with by using the mean from answered items only.

PFDI-20 Summary Score: Add the scores from the 3 scales together to obtain the summary score (range 0 to 300).

Continued on following page

PFIQ – 7 Instructions: Some women find that bladder, bowel, or vaginal symptoms affect their activities, relationships, and feelings. For each question place an check mark in the response that best describes how much your activities, relationships, or feelings have been affected by your bladder, bowel, or vaginal symptoms or conditions **over the last 3 months**. Please make sure you mark an answer in all 3 columns for each question.

How do symptoms or conditions relating to the following → → → Usually affect your ...↓	Bladder or urine	Bowel or rectum	Vagina or pelvis
1. Ability to do household chores (cooking, housecleaning, laundry)?	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit
2. Ability to do physical activities such as walking, swimming, or other exercise?	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit
3. Entertainment activities such as going to a movie or concert?	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit
4. Ability to travel by car or bus for a distance greater than 30 minutes away from home?	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit
5. Participating in social activities outside your home?	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit
6. Emotional health (nervousness, depression, etc.)?	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit
7. Feeling frustrated?	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit

All of the items use the following response scale:
0 = not at all; 1 = somewhat, 2 = moderately, 3 = quite a bit

Scales:

Urinary Impact Questionnaire (UIQ-7); 7 items under column heading “Bladder or urine.” Colorectal-Anal Impact Questionnaire (CRAIQ-7): 7 items under column heading “Bowel or rectum.”
Pelvic Organ Prolapse Impact Questionnaire (POPIQ-7): 7 items under column heading “Pelvis or vagina.”

Scale scores: Obtain the mean value for all of the answered items within the corresponding scale (possible value 0 to 3) and then multiply by 100/3) to obtain the scale score (range 0 to 100). Missing items are dealt with by using the mean from answered items only.

Total score of each section _____ divided by 7 _____ X 33.3 = _____

PFIQ-7 Summary Score: Add the scores from the 3 scales together to obtain the summary score (range 0 to 300).

Barber, M., Walters, M., et al. (2005). "Short forms of two condition-specific quality of life questionnaires for women with pelvic floor disorders (PFDI-20 and PFIQ -7)." American Journal of Obstetrics and Gynecology 193: 103-113.